

Exact calculations of vertex $\bar{s}\gamma b$ and $\bar{s}Zb$ in the unitary gauge

Lu-sen Wu and Zhen-jun Xiao*

*Department of Physics and Institute of Theoretical Physics,
Nanjing Normal University, Nanjing, Jiangsu 210097, P.R.China*

(Dated: February 2, 2008)

Abstract

In this paper, we present the exact calculations for the vertex $\bar{s}\gamma b$ and $\bar{s}Zb$ in the unitary gauge. We found that (a) the divergent- and μ -dependent terms are left in the effective vertex function $\Gamma_\mu^\gamma(p, k)$ for $b \rightarrow s\gamma$ transition even after we sum up the contributions from four related Feynman diagrams; (b) for an on-shell photon, such terms do not contribute et al; (c) for off-shell photon, these terms will be canceled when the contributions from both vertex $\bar{s}\gamma b$ and $\bar{s}Zb$ are taken into account simultaneously, and therefore the finite and gauge independent function $Z_0(x_t) = C_0(x_t) + D_0(x_t)/4$, which governs the semi-leptonic decay $b \rightarrow sl^-l^+$, is derived in the unitary gauge.

PACS numbers: 13.25.Hw, 12.38.Bx, 12.20.Ds, 11.15.Bt

*Electronic address: xiaozhenjun@njnu.edu.cn

I. INTRODUCTION

The inclusive rare radiative decays of b quark, $b \rightarrow sV$ with $V = \gamma, Z$ followed by the decays $V \rightarrow q\bar{q}, l^+l^-$, are induced by the flavor changing neutral current (FCNC) and play an important role in testing the standard model (SM) and searching for the signal or evidence of the new physics beyond the SM [1].

Because of its great importance, these rare decays have been studied extensively during the past three decades. The $s \rightarrow dV$ transitions with $V = \gamma, Z$ were firstly calculated in Ref. [2] in R_ξ gauge. The general transition $q_i \rightarrow q_j\gamma$ for arbitrary quark flavors with arbitrary mass of internal quarks were evaluated in the 't Hooft-Feynman gauge [3] or in a non-linear R_ξ gauge [4], where the current conservation was used to renormalize the vertex $\bar{q}_j\gamma q_i$.

In Ref. [5], an exact calculation of the $\bar{d}s\gamma$ vertex was made in the unitary gauge, where the Ward-Takahashi identity was used to renormalize the vertex. For the quark-photon vertex $\bar{d}s\gamma$ in the unitary gauge, it was firstly calculated in Ref. [6], but the results given there were not correct. In Ref. [7], the author studied the radiative decay $b \rightarrow s\gamma$ directly and presented their new results for the on-shell vertex function $\Gamma_\mu^\gamma(p, k)$. The analytical results presented in Refs. [2, 3, 4] can also be extended easily to the case of vertex $\bar{s}\gamma b$, the relevant basic functions $D_0(x_t)$ and $D'_0(x_t)$ [8, 9] can be extracted directly. Of course, it is worth of mentioning that the basic function $D_0(x_t)$, as well as $B_0(x_t)$ and $C_0(x_t)$ are all gauge dependent [8, 9].

In this paper, we present the results of recalculation of the $\bar{s}\gamma b$ vertex and the first calculation of $\bar{s}Zb$ vertex in the unitary gauge. This paper is organized as follows. In Sec. II, we recalculate the $\bar{s}\gamma b$ vertex in the unitary gauge and extract out the basic function $D_0(x_t)$ and $D'_0(x_t)$. In Sec. III, we calculate the $\bar{s}Zb$ vertex in the unitary gauge, derive the basic function $C_0(x_t)$ and make a linear combination of $Z_0(x_t) = C_0(x_t) + D_0(x_t)/4$. The summary and some discussions are also included in Sec. III.

II. THE EFFECTIVE $\bar{s}\gamma b$ COUPLING

In the unitary gauge, the one loop Feynman diagrams for the flavor changing vertex $\Gamma_\mu^{(\gamma)}(p, k)$ are illustrated in Fig. 1 for the case of $V = \gamma$.

For Fig. 1a, with the aid of the unitarity relation $\sum_{j=u,c,t} V_{js}^* V_{jb} = 0$, the vertex $\Gamma_\mu^a(p, k)$ can be written as

$$\begin{aligned} \Gamma_\mu^a(p, k) = & \frac{2}{3}e\frac{g^2}{32\pi^2} \sum_j V_{js}^* V_{jb} \cdot \left\{ \left[-\frac{3}{4-N} + \frac{3}{2} \ln \left(\frac{e^\gamma M_W^2}{4\pi\mu^2} \right) - \frac{5}{6} \right] x_j \gamma_\mu L + E_1 \gamma_\mu L \right. \\ & + (E_2 \not{p} \gamma_\mu \not{p} + E_3 \not{k} \gamma_\mu \not{k} + E_4 \not{p} \gamma_\mu \not{k} + E_5 \not{k} \gamma_\mu \not{p} + E_6 \not{p} \not{k} \gamma_\mu + E_7 \gamma_\mu \not{k} \not{p}) \frac{L}{M_W^2} \\ & \left. + E_8 \not{p} \not{k} \gamma_\mu \not{k} \not{p} \frac{L}{M_W^4} \right\}, \end{aligned} \quad (1)$$

where $x_j = m_j^2/M_W^2$ with $m_j = (m_u, m_c, m_t)$, $L = (1 - \gamma_5)/2$, $N = 4 - 2\epsilon$ is the dimension of space-time in dimensional regularization, and the functions

$$E_i = \int_0^1 dx \int_0^{1-x} dy H_i, \quad i = 1, 2, 3, \dots, 8. \quad (2)$$

with

$$\begin{aligned}
H_1 &= \left\{ 2x_j + \frac{[x(1-x)p^2 + y(1-y)k^2 - 2xyp \cdot k][x(1-x)p^2 - y^2k^2 - 2xyp \cdot k]}{M_W^4} \right\} D^{-1} \\
&\quad - \left[2(1-3x) - (5-6x)x_j + \frac{12x(1-x)p^2 + 3y(3-4y)k^2 - 24xyp \cdot k}{M_W^2} \right] \ln D, \\
H_2 &= [2x^2 + (1-x)^2x_j] D^{-1} - \ln D, \\
H_3 &= [-2y(1-y) + y^2x_j] D^{-1}, \\
H_4 &= -[2x(1-y) + (1-x)yx_j] D^{-1} + \ln D, \\
H_5 &= [2xy - (1-x)yx_j] D^{-1}, \\
H_6 &= -(1-x-y) \left\{ \frac{1}{M_W^2} [x(1-x)p^2 - y^2k^2 - 2xyp \cdot k] D^{-1} - 3 \ln D \right\}, \\
H_7 &= y \left\{ \frac{1}{M_W^2} [x(1-x)p^2 + y(1-y)k^2 - 2xyp \cdot k] D^{-1} - 3 \ln D \right\}, \\
H_8 &= -y(1-x-y)D^{-1}, \tag{3}
\end{aligned}$$

with

$$D = x + (1-x)x_j - \frac{1}{M_W^2}[x(1-x)p^2 + y(1-y)k^2 - 2xy(p \cdot k)]. \tag{4}$$

Here the functions E_i and H_i are identical with those presented in Ref. [5].

When the terms proportional to p^2/M_W^2 , k^2/M_W^2 and $p \cdot k/M_W^2$ in the functions H_i and D are neglected, it is easy to do the integration and find the analytical expressions of function E_i . From the analytical results, we confirmed that the vertex function $\Gamma_\mu^a(p, k)$ in Eq. (1) are identical with those presented in Ref. [5] for the $\bar{d}sg$ vertex, after making a proper change of the quarks involved and a replacement of the factor $g_s \lambda^a/2$ to eQ_u ($Q_u = 2/3$ is the charge fraction of the up-type quarks).

For $b \rightarrow sg$ decay, Fig. 1b is irrelevant. For $b \rightarrow s\gamma$ decay, however, Fig. 1b must be taken into account. For Fig. 1b, the decay amplitude is of the form

$$\begin{aligned}
\mathcal{M}_{1b} &= \bar{u}_s(p') \left\{ \int \frac{d^4q}{(2\pi)^4} \sum_j \frac{ig}{\sqrt{2}} \gamma_\alpha LV_{js}^* \frac{i}{\not{p} - \not{q} - m_j} \frac{ig}{\sqrt{2}} \gamma_\beta LV_{jb} \right. \\
&\quad \cdot \frac{i}{(k-q)^2 - M_W^2} \left[-g^{\alpha\lambda} + \frac{1}{M_W^2} (k^\alpha - q^\alpha)(k^\lambda - q^\lambda) \right] \\
&\quad \cdot ie[(2q_\mu - k_\mu)g_{\nu\lambda} + (2k_\nu - q_\nu)g_{\lambda\mu} + (-k_\lambda - q_\lambda)g_{\mu\nu}] \\
&\quad \cdot \frac{i}{q^2 - M_W^2} \left[-g^{\beta\nu} + \frac{q^\beta q^\nu}{M_W^2} \right] \left. \right\} u_b(p) \epsilon^\mu(k). \tag{5}
\end{aligned}$$

The corresponding vertex $\Gamma_\mu^b(p, k)$ can then be written as

$$\Gamma_\mu^b(p, k) = ie \frac{g^2}{2} \sum_j V_{js}^* V_{jb} \int \frac{d^4q}{(2\pi)^4} \frac{N_\mu L}{[(p-q)^2 - m_j^2][(k-q)^2 - M_W^2][q^2 - M_W^2]}, \tag{6}$$




FIG. 1: The One-loop Feynman diagrams contributing to the induced $\bar{s}\gamma b$ vertex or $\bar{s}Zb$ vertex in the unitary gauge.

where the numerator N_μ is of the form

$$\begin{aligned}
N_\mu = & \gamma_\alpha (\not{p} - \not{q}) \gamma^\alpha (2q_\mu - k_\mu) + \gamma_\mu (\not{p} - \not{q}) (2\not{k} - \not{q}) - (\not{k} + \not{q}) (\not{p} - \not{q}) \gamma_\mu \\
& + \frac{1}{M_W^2} [(\not{k} - \not{q}) (\not{p} - \not{q}) \not{q} q_\mu - (\not{k} - \not{q}) (\not{p} - \not{q}) q^2 \gamma_\mu + \dots] \\
& + \frac{1}{M_W^4} (\not{k} - \not{q}) (\not{p} - \not{q}) \not{q} (k \cdot q k_\mu - k^2 q_\mu).
\end{aligned} \tag{7}$$

By employing the Feynman parametrization and dimensional regularization, we find the expression of the vertex $\Gamma_\mu^b(p, k)$

$$\begin{aligned}
\Gamma_\mu^b(p, k) = & e \frac{g^2}{32\pi^2} \sum_j V_{js}^* V_{jb} \left\{ \left[\frac{3}{4-N} - \frac{3}{2} \ln \left(\frac{e^\gamma M_W^2}{4\pi\mu^2} \right) \right] x_j \gamma_\mu L \right. \\
& + \left[\frac{1}{2} \frac{1}{4-N} - \frac{1}{4} \ln \left(\frac{e^\gamma M_W^2}{4\pi\mu^2} \right) + \frac{1}{6} \right] x_j (k^2 \gamma_\mu - \not{k} k_\mu) \frac{L}{M_W^2} \\
& + E'_1 \gamma_\mu L + [E'_2 \not{p} \gamma_\mu \not{p} + E'_3 \not{k} \gamma_\mu \not{k} + E'_4 \not{p} \gamma_\mu \not{k} \\
& \left. + E'_5 \not{k} \gamma_\mu \not{p} + E'_6 \not{p} \not{k} \gamma_\mu + E'_7 \gamma_\mu \not{k} \not{p}] \frac{L}{M_W^2} \right\},
\end{aligned} \tag{8}$$

where the functions E'_i have been defined as

$$E'_i = \int_0^1 dx \int_0^{1-x} dy H'_i, \quad i = 1, 2, 3, \dots, 7. \tag{9}$$

with

$$\begin{aligned}
H'_1 = & \left\{ \frac{1}{M_W^2} [-4x(1-x)p^2 + 2(1-x-2y+4xy)p \cdot k - 2y(1-2y)k^2] \right. \\
& + \frac{1}{M_W^4} \left[x^3(1-x)p^4 - \frac{1}{2}x(1-x+2y-2y^2-3xy+4xy^2)p^2k^2 \right. \\
& - 4xy(1-x)(1-y)(p \cdot k)^2 - 2x^2(1-x)(1-2y)p^2(p \cdot k) \\
& + (1-x-y-3y^2+2y^3+xy+5xy^2-4xy^3)k^2(p \cdot k) - \frac{1}{2}y(1-y-3y^2+2y^3)k^4 \left. \right] \\
& + \frac{1}{M_W^6} \left[\frac{1}{2}x^3(1-x)p^4k^2 - \frac{1}{2}x^2(1-x-2y+3xy)p^2k^2(p \cdot k) + \frac{1}{2}xy(1-y-xy)p^2k^4 \right. \\
& - xy(1-x)(1-y)k^2(p \cdot k)^2 + \frac{1}{2}xy^2(1-y)k^4(p \cdot k) \left. \right] \left. \right\} D'^{-1} \\
& + \left\{ 6 + \frac{1}{M_W^2} \left[x(5-7x)p^2 - 2(2-3x-4y+7xy)(p \cdot k) + (1 + \frac{9}{2}y - 7y^2)k^2 \right] \right. \\
& + \frac{1}{M_W^4} \left[-\frac{9}{4}x^2p^2k^2 - \frac{1}{2}(1-3x-2y+5xy)k^2(p \cdot k) - \frac{1}{4}y^2k^4 \right] \left. \right\} \ln D' \\
& - 9D' \ln D' - \frac{3}{4} \frac{k^2}{M_W^2} D' \ln D', \tag{10}
\end{aligned}$$

$$\begin{aligned}
H'_2 = & \left\{ -2x(1-x) \right. \\
& - \left(\frac{1}{M_W^2} - \frac{k^2}{2M_W^4} \right) [x^3(1-x)p^2 - 2x^3y(p \cdot k) + xy(1-y-xy)k^2] \left. \right\} D'^{-1} \\
& + x(1+4x) \left(1 - \frac{1}{2} \frac{k^2}{M_W^2} \right) \ln D', \tag{11}
\end{aligned}$$

$$\begin{aligned}
H'_3 = & \left\{ -y(1-2y) + \frac{1}{2} \frac{1}{M_W^2} [x(1-x+4y-3xy-4y^2+2xy^2)p^2 \right. \\
& - 2(1-x)(1-y-3y^2+2y^3)(p \cdot k) + y(1-y-3y^2+2y^3)k^2] \\
& + \frac{1}{2} \frac{p \cdot k}{M_W^4} [x^2(1-x-2y+xy)p^2 + 2xy(1-x-y+xy)(p \cdot k) - xy^2(1-y)k^2] \left. \right\} D'^{-1} \\
& + \left\{ \left(-1 - \frac{9}{2}y + 4y^2 \right) + \frac{1}{2} \frac{1}{M_W^2} \left[-x \left(1 - \frac{1}{2}x \right) p^2 \right. \right. \\
& + (1-3x-2y+5xy)(p \cdot k) + \frac{1}{2}y^2k^2 \left. \right] \left. \right\} \ln D' + \frac{3}{4} D' \ln D', \tag{12}
\end{aligned}$$

$$\begin{aligned}
H'_4 = & \left\{ (1-x-2y+2xy) + \frac{1}{2} \frac{1}{M_W^2} [x^2(1-x-2y+2xy)p^2 - 2x^2y(1-2y)(p \cdot k) \right. \\
& + (1-x-y-3y^2-xy^2+2y^3+2xy^3)k^2] \\
& + \frac{1}{2} \frac{p \cdot k}{M_W^4} [-x^3(1-x)p^2 + 2x^3y(p \cdot k) - xy(1-y-xy)k^2] \left. \right\} D'^{-1} \\
& + \frac{1}{2} \left\{ (-1-3x+2y+8xy) + \frac{1}{M_W^2} \left[(x+4x^2)(p \cdot k) - \frac{k^2}{2}(1-2y) \right] \right\} \ln D', \quad (13)
\end{aligned}$$

$$\begin{aligned}
H'_5 = & \left\{ 2xy + \left(\frac{1}{M_W^2} - \frac{1}{2} \frac{k^2}{M_W^4} \right) [x^2(1-x-2y+xy)p^2 \right. \\
& + 2xy(1-x)(1-y)(p \cdot k) - xy^2(1-y)k^2] \left. \right\} D'^{-1} \\
& + (-3x+4xy) \left(1 - \frac{1}{2} \frac{k^2}{M_W^2} \right) \ln D', \\
H'_6 = & \left\{ (2-2x-y) + \frac{1}{2} \frac{1}{M_W^2} [x^2(1-x)p^2 - 2x^2y(p \cdot k) \right. \\
& + (-1+x+y-y^2-xy^2)k^2] + \frac{1}{2} \frac{1}{M_W^4} [x^2(1-x-2y+xy)p^2k^2 + 2x^3y(p \cdot k)^2 \\
& - x^3(1-x)p^2(p \cdot k) + xy(1-2x-y+3xy)k^2(p \cdot k) - xy^2(1-y)k^4] \left. \right\} D'^{-1} \\
& + \left\{ \frac{1}{2}(1-3x-2y) + \frac{1}{M_W^2} \left[\frac{1}{2}x(1+4x)(p \cdot k) + \frac{1}{4}(1-6x+8xy)k^2 \right] \right\} \ln D', \\
H'_7 = & \left\{ (-2+2x+y) - \frac{1}{M_W^2} [x^2yp^2 - 2xy(1-y)(p \cdot k) - y^2(2-y)k^2] \right\} D'^{-1} \\
& + \left[(1-2y) - \frac{1}{2}y \frac{k^2}{M_W^2} \right] \ln D', \quad (14)
\end{aligned}$$

and

$$D' = 1 - x + xx_j - \frac{1}{M_W^2} [x(1-x)p^2 + y(1-y)k^2 - 2xyp \cdot k]. \quad (15)$$

For Fig. 1c and 1d, we find the vertex $\Gamma_\mu^{(c+d)}(p, k)$ in the same way

$$\begin{aligned}
\Gamma_\mu^{(c+d)}(p, k) = & e \frac{g^2}{32\pi^2} \sum_j V_{js}^* V_{jb} \left\{ \left[-\frac{1}{4-N} + \frac{1}{2} \ln \left(\frac{e^\gamma M_W^2}{4\pi\mu^2} \right) - \frac{2}{9} \right] x_j \gamma_\mu L \right. \\
& + \frac{1}{3} \frac{m_b^2 F(m_b^2) - m_s^2 F(m_s^2)}{m_b^2 - m_s^2} \gamma_\mu L \\
& \left. + \frac{1}{3} \frac{m_b m_s (F(m_b^2) - F(m_s^2))}{m_b^2 - m_s^2} \gamma^\mu R \right\}, \quad (16)
\end{aligned}$$

where $(L, R) = (1 \mp \gamma_5)/2$, and the function $F(p^2)$ is

$$\begin{aligned}
F(p^2) = & \int_0^1 dx \left[x(2-3x) + (1-x)(4-3x)x_j - x(1-x)(5-4x) \frac{p^2}{M_W^2} \right] \\
& \cdot \ln \left[x + (1-x)x_j - x(1-x) \frac{p^2}{M_W^2} \right]. \quad (17)
\end{aligned}$$

By combining the four pieces together, we find the total vertex function $\Gamma_\mu^\gamma(p, k)$

$$\begin{aligned}
\Gamma_\mu^\gamma(p, k) &= \Gamma_\mu^a(p, k) + \Gamma_\mu^b(p, k) + \Gamma_\mu^{(c+d)}(p, k) \\
&= e \frac{g^2}{32\pi^2} \sum_j V_{js}^* V_{jb} \left[\frac{1}{2} \frac{1}{4-N} - \frac{1}{4} \ln \left(\frac{e^\gamma M_W^2}{4\pi\mu^2} \right) + \frac{1}{6} \right] x_j (k^2 \gamma_\mu - \not{k} k_\mu) \frac{L}{M_W^2} \\
&\quad + e \frac{g^2}{32\pi^2} \sum_j V_{js}^* V_{jb} \left\{ -\frac{7}{9} x_j \gamma_\mu L + \frac{1}{3} \frac{m_b^2 F(m_b^2) - m_s^2 F(m_s^2)}{m_b^2 - m_s^2} \gamma_\mu L \right. \\
&\quad + \frac{1}{3} \frac{m_b m_s (F(m_b^2) - F(m_s^2))}{m_b^2 - m_s^2} \gamma_\mu R + \tilde{E}_1 \gamma_\mu L \\
&\quad + \left(\tilde{E}_2 \not{p} \gamma_\mu \not{p} + \tilde{E}_3 \not{k} \gamma_\mu \not{k} + \tilde{E}_4 \not{p} \gamma_\mu \not{k} + \tilde{E}_5 \not{k} \gamma_\mu \not{p} + \tilde{E}_6 \not{p} \not{k} \gamma_\mu + \tilde{E}_7 \gamma_\mu \not{k} \not{p} \right) \frac{L}{M_W^2} \\
&\quad \left. + \tilde{E}_8 \not{p} \not{k} \gamma_\mu \not{k} \not{p} \frac{L}{M_W^4} \right\} \tag{18}
\end{aligned}$$

where

$$\tilde{E}_i = \frac{2}{3} E_i + E'_i, \quad i = 1, 2, 3, \dots, 7; \tag{19}$$

$$\tilde{E}_8 = \frac{2}{3} E_8. \tag{20}$$

Following Ref. [2], the renormalization of the vertex $\bar{s}\gamma b$ is achieved here by summing up the contributions from all four Feynman diagrams in Fig. 1. For a given process, in principle, the divergent terms from individual Feynman diagrams should be canceled each other when combining all contributions. In fact, the divergent term and the μ -dependent term proportional to $x_j \gamma_\mu L$ in Eqs. (1,8) and (16) do cancel each other, but the terms in the square brackets proportional to $(k^2 \gamma_\mu - \not{k} k_\mu)$ in Eq. (8), i.e.,

$$T^{div} = \frac{1}{2} \frac{1}{4-N} - \frac{1}{4} \ln \left(\frac{e^\gamma M_W^2}{4\pi\mu^2} \right) \tag{21}$$

are still left in the total vertex function $\Gamma_\mu^\gamma(p, k)$. The term T^{div} arises from Fig. 1b, the terms proportional to $1/M_W^4$ in Eq. (7).

If we use the Ward-Takahashi identity directly to renormalize the vertex, as being done in Refs. [5, 6], we find the same vertex function $\Gamma_\mu^\gamma(p, k)$ as in Eq. (18).

Based on previous studies [2, 9], we know that the effective $\bar{s}\gamma b$ coupling can be generally written as

$$\begin{aligned}
\bar{s}\gamma b &= i \bar{s} \Gamma_\mu^\gamma(p, k) b = i \sum_j V_{js}^* V_{jb} \frac{G_F}{\sqrt{2}} \frac{e}{8\pi^2} \{ D_0(x_j) \bar{s} (k^2 \gamma_\mu - \not{k} k_\mu) (1 - \gamma_5) b \\
&\quad - i D'_0(x_j) \bar{s} \sigma_{\mu\nu} k^\nu [m_s (1 - \gamma_5) + m_b (1 + \gamma_5)] b \}, . \tag{22}
\end{aligned}$$

where the basic functions $D_0(x_j)$ and $D'_0(x_j)$ were obtained many years ago [2]¹ and takes

¹ In Ref. [2], the terms $F_1(x_j)$ and $F_2(x_j)$ instead of $D_0(x_j)$ and $D'_0(x_j)$ were used to denote the two form factors.

the following form in the 't Hooft-Feynman gauge [2, 8]

$$D_0(x_t) = \frac{-25x_t^2 + 19x_t^3}{36(1-x_t)^3} - \frac{8 - 32x_t + 54x_t^2 - 30x_t^3 + 3x_t^4}{18(1-x_t)^4} \ln[x_t], \quad (23)$$

$$D'_0(x_t) = -\frac{8x_t^3 + 5x_t^2 - 7x_t}{12(1-x_t)^3} + \frac{x_t^2(2-3x_t)}{2(1-x_t)^4} \ln[x_t], \quad (24)$$

where $x_t = m_t^2/M_W^2$.

In order to compare our results with those given in Eqs. (23) and (24), we do the integration and extract out the corresponding basic function $D_0(x_t)$ and $D'_0(x_t)$ from vertex function $\Gamma_\mu^\gamma(p, k)$ in Eq. (18). Since m_b^2, m_s^2, k^2 are all much smaller than M_W^2 , the terms proportional to $p^2/M_W^2, k^2/M_W^2$ and $k \cdot p/M_W^2$ in functions H_i and H'_i can be neglected safely. The x - and y -integrations in Eq.(20) can therefore be evaluated easily. We then find the analytical expressions for the functions $D_0(x_t)$ and $D'_0(x_t)$ in the unitary gauge:

$$D_0(x_t) = T^{div} + \frac{153x_t - 383x_t^2 + 245x_t^3 - 27x_t^4}{72(1-x_t)^3} - \frac{16 - 64x_t + 36x_t^2 + 93x_t^3 - 84x_t^4 + 9x_t^5}{36(1-x_t)^4} \ln[x_t], \quad (25)$$

$$D'_0(x_t) = -\frac{8x_t^3 + 5x_t^2 - 7x_t}{12(1-x_t)^3} + \frac{x_t^2(2-3x_t)}{2(1-x_t)^4} \ln[x_t], \quad (26)$$

where $x_t = m_t^2/M_W^2$ and the term T^{div} has been defined in Eq. (21). It is easy to see that

- The basic function $D'_0(x_t)$ in both 't Hooft-Feynman and unitary gauge are identical, since $D'_0(x_t)$ itself is gauge independent[8, 9];
- The basic function $D_0(x_t)$ in Eq. (25) is different from that in the 't Hooft-Feynman gauge. The reason is that the basic function $D_0(x_t)$, as well as $B_0(x_t)$ and $C_0(x_t)$ are all gauge-dependent [8]. The basic function $B_0(x_t)$ and $C_0(x_t)$ come from the box diagrams of $B^0 - \bar{B}^0$ mixing and from the Z -penguin diagrams with internal top quark propagators, respectively.

For the case of a real photon emission, the photon is on-shell which means $k^2 = 0$, and the k_μ term can not contribute either by using the transverse polarization condition $k \cdot \epsilon = 0$. For a on-shell γ , therefore, only the term proportional to $D'_0(x_t)$ in Eq. (22) contribute, the physics is consequently independent of the gauge. For a time-like photon ($k^2 > 0$), the divergent term T^{div} may induce a serious problem! But for an off-shell γ , the photon will interact with other particles, new Feynman diagrams will be involved and need to be sum up, which should lead to a finite and gauge independent final result.

For the inclusive $b \rightarrow sl^-l^+$ decays, for example, the contributions from both decay chains $b \rightarrow s\gamma^* \rightarrow sl^-l^+$ and $b \rightarrow sZ^* \rightarrow sl^-l^+$ should be considered simultaneously. We indeed find that the term T^{div} in Eq. (21) is canceled by the corresponding term in the vertex $\Gamma_\mu^Z(p, k)$ when the contribution from the decay chain $b \rightarrow sZ^* \rightarrow sl^-l^+$ was also taken into account. We will show this point explicitly in next section.

As mentioned in the introduction, the Fig. 1b was firstly calculated in the unitary gauge in Ref. [6], but the analytical results given there were wrong. In Ref. [7], the author

studied the radiative decay $b \rightarrow s\gamma$ directly in the unitary gauge and presented their new results for the on-shell vertex function $\Gamma_\mu^\gamma(p, k)$:

$$\begin{aligned} \Gamma_\mu^\gamma(p, k) = & \frac{eG_F}{4\sqrt{2}\pi^2} \sum_j V_{js}^* V_{jb} [A_{tot}(x_j) (k_\mu \not{k} - k^2 \gamma_\mu) L \\ & + iB_{tot}(x_j) \sigma_{\mu\nu} k^\nu (m_s L + m_b R)], \end{aligned} \quad (27)$$

where $L, R = (1 \mp \gamma_5)/2$, and

$$A_{tot}(x_t) = -D_0(x_t), \quad (28)$$

$$B_{tot}(x_t) = D'_0(x_t), \quad (29)$$

by definition. Using the formulae as given in Eqs.(6-13) of ref. [7], one finds the analytical expressions of $D_0(x_t)$ and $D'_0(x_t)$

$$\begin{aligned} D_0(x_t) = -A_{tot}(x_t) = & \frac{63x_t - 151x_t^2 + 82x_t^3}{36(1-x_t)^3} \\ & - \frac{16 - 127x_t + 234x_t^2 - 123x_t^3 + 6x_t^4}{36(1-x_t)^4}, \end{aligned} \quad (30)$$

$$D'_0(x_t) = B_{tot}(x_t) = -\frac{8x_t^3 + 5x_t^2 - 7x_t}{12(1-x_t)^3} + \frac{x_t^2(2-3x_t)}{2(1-x_t)^4} \ln[x_t]. \quad (31)$$

It is easy to see that

- The gauge independent function $D'_0(x_t)$ as given in Eqs.(24,26) and (31) are identical.
- For function $D_0(x_t)$, however, the two expressions in Eqs.(25) and (30) are different.

In Ref. [8], the gauge independent function $Z_0(x_t)$, which governs the semi-leptonic decay $b \rightarrow sl^+l^-$, was defined as a linear combination of $C_0(x_t, \xi)$ and $D_0(x_t, \xi)$,

$$\begin{aligned} Z_0(x_t) = & C_0(x_t, \xi) + \frac{1}{4}D_0(x_t, \xi) \\ = & \frac{108x_t - 259x_t^2 + 163x_t^3 - 18x_t^4}{144(1-x_t)^3} - \frac{8 - 50x_t + 63x_t^2 + 6x_t^3 - 24x_t^4}{72(1-x_t)^4} \ln[x_t], \end{aligned} \quad (32)$$

where

$$C_0(x_t, \xi) = C_0(x_t) + \frac{1}{2}\bar{\rho}(x, \xi), \quad D_0(x_t, \xi) = D_0(x_t) - 2\bar{\rho}(x, \xi), \quad (33)$$

with

$$\bar{\rho}(x, \xi) = 2\rho(x, \xi) - 7B_0(x_t), \quad (34)$$

$$\begin{aligned} \rho(x, \xi) = & \frac{\xi}{x-\xi} \left(\frac{3}{4} \frac{1}{x-1} \right) x \ln[x] + \frac{1}{8} \frac{\xi^2}{x-\xi} \left[\left(\frac{5+\xi}{1-\xi} - \frac{\xi}{x-\xi} \right) \ln[\xi] - 1 \right] \\ & - (x \rightarrow 0), \end{aligned} \quad (35)$$

$$B_0(x_t) = \frac{x_t}{4(1-x_t)} + \frac{x_t}{4(1-x_t)^2} \ln[x-t]. \quad (36)$$

In next section, we will calculate the Z -penguin diagrams, extract out the basic function $C_0(x_t)$, and check the gauge independence of function $Z_0(x_t) = C_0(x_t) + D_0(x_t)/4$ in the unitary gauge.

III. THE EFFECTIVE $\bar{s}Zb$ COUPLING

In the unitary gauge, the one loop Feynman diagrams for the flavor changing vertex $\Gamma_\mu^Z(p, k)$ are illustrated in Fig. 1 for the case of $V = Z$ with a virtual Z ($k^2 > 0$).

Following the same procedure as in last section, one can find the vertex functions from Fig. 1a-1d for the case of $V = Z$. First, the individual decay amplitudes can be written as

$$\begin{aligned} \mathcal{M}_Z^{(a)} = & \bar{u}_s(p') \left\{ \int \frac{d^4 q}{(2\pi)^4} \sum_j \frac{ig}{\sqrt{2}} \gamma_\alpha L V_{js}^* \frac{i}{\not{p} - \not{q} - \not{k} - m_j} \right. \\ & \cdot \frac{ig}{4 \cos \theta} \gamma_\mu \left[\left(1 - \frac{8}{3} \sin^2 \theta\right) - \gamma_5 \right] \frac{i}{\not{p} - \not{q} - m_j} \\ & \cdot \left. \frac{ig}{\sqrt{2}} \gamma_\beta L V_{jb} \frac{i}{q^2 - M_W^2} \left(-g^{\alpha\beta} + \frac{q^\alpha q^\beta}{M_W^2} \right) \right\} u_b(p) \epsilon^\mu(k), \end{aligned} \quad (37)$$

$$\begin{aligned} \mathcal{M}_Z^{(b)} = & \bar{u}_s(p') \left\{ \int \frac{d^4 q}{(2\pi)^4} \sum_j \frac{ig}{\sqrt{2}} \gamma_\alpha L V_{js}^* \frac{i}{\not{p} - \not{q} - m_j} \frac{ig}{\sqrt{2}} \gamma_\beta L V_{jb} \right. \\ & \cdot \frac{i}{(k - q)^2 - M_W^2} \left[-g^{\alpha\lambda} + \frac{1}{M_W^2} (k^\alpha - q^\alpha)(k^\lambda - q^\lambda) \right] \\ & \cdot ig \cos \theta [(2q_\mu - k_\mu)g_{\nu\lambda} + (2k_\nu - q_\nu)g_{\lambda\mu} + (-k_\lambda - q_\lambda)g_{\mu\nu}] \\ & \cdot \left. \frac{i}{q^2 - M_W^2} \left[-g^{\beta\nu} + \frac{q^\beta q^\nu}{M_W^2} \right] \right\} u_b(p) \epsilon^\mu(k), \end{aligned} \quad (38)$$

$$\begin{aligned} \mathcal{M}_Z^{(c+d)} = & \bar{u}_s(p') \left\{ -\frac{ig}{2 \cos \theta} \cdot \frac{g^2}{32\pi^2} \sum_j V_{js}^* V_{jb} \left[\left(-1 + \frac{2}{3} \sin^2 \theta\right) \left(-\frac{3}{4 - N}\right) \right. \right. \\ & + \frac{3}{2} \ln \left(\frac{e^\gamma M_W^2}{4\pi\mu^2} \right) - \frac{2}{3} \Big] x_j \gamma_\mu L + \left(-1 + \frac{2}{3} \sin^2 \theta\right) \frac{m_b^2 F(m_b^2) - m_s^2 F(m_s^2)}{m_b^2 - m_s^2} \gamma_\mu L \\ & \left. + \frac{2}{3} \sin^2 \theta \frac{m_b m_s (F(m_b^2) - F(m_s^2))}{m_b^2 - m_s^2} \gamma_\mu R \right] \Big\} u_b(p) \epsilon^\mu(k). \end{aligned} \quad (39)$$

Taking the same approximation as in last section, the corresponding vertex function $\Gamma_\mu^Z(p, k)$ can be derived from Eqs. (37)-(39) after making the x - and y -integrations:

$$\begin{aligned} \Gamma_\mu^{(Z,a)}(p, k) = & \frac{g}{2 \cos \theta} \frac{g^2}{32\pi^2} \sum_j V_{js}^* V_{jb} \left\{ \left(1 - \frac{4}{3} \sin^2 \theta\right) \left[-\frac{3}{4 - N} + \frac{3}{2} \ln \left(\frac{e^\gamma M_W^2}{4\pi\mu^2} \right) - \frac{5}{6} \right] x_j \right. \\ & + \left(1 - \frac{4}{3} \sin^2 \theta\right) \left[\frac{8 - 31x_j + 5x_j^2}{12(1 - x_j)} - \frac{3x_j^2(2 - x_j)}{2(1 - x_j)^2} \ln x_j \right] \\ & - \left[\frac{1}{4 - N} - \frac{1}{2} \ln \left(\frac{e^\gamma M_W^2}{4\pi\mu^2} \right) - \frac{1}{2} \right] x_j \\ & \left. + \frac{x_j(5 + x_j)}{4(1 - x_j)} + \frac{x_j(x_j^2 - 2x_j + 4)}{2(1 - x_j)^2} \ln x_j \right\} \gamma_\mu L + \dots, \end{aligned} \quad (40)$$

$$\begin{aligned}\Gamma_\mu^{(Z,b)}(p,k) = & g \cos \theta \frac{g^2}{32\pi^2} \sum_j V_{js}^* V_{jb} \left\{ \left[\frac{3}{4-N} - \frac{3}{2} \ln \left(\frac{e^\gamma M_W^2}{4\pi\mu^2} \right) \right] x_j \right. \\ & \left. + \frac{-2+13x_j-5x_j^2}{4(1-x_j)} + \frac{3x_j^2(2-x_j)}{2(1-x_j)^2} \ln x_j \right\} \gamma_\mu L + \dots, \quad (41)\end{aligned}$$

and

$$\begin{aligned}\Gamma_\mu^{(Z,c+d)}(p,k) = & \frac{g}{2 \cos \theta} \left(1 - \frac{2}{3} \sin^2 \theta \right) \frac{g^2}{32\pi^2} \sum_j V_{js}^* V_{jb} \\ & \cdot \left\{ \left[-\frac{3}{4-N} + \frac{3}{2} \ln \left(\frac{e^\gamma M_W^2}{4\pi\mu^2} \right) - \frac{2}{3} \right] x_j \right. \\ & \left. - \frac{2+23x_j-7x_j^2}{12(1-x_j)} - \frac{3x_j^2(2-x_j)}{2(1-x_j)^2} \ln x_j \right\} \gamma_\mu L + \dots. \quad (42)\end{aligned}$$

where only the terms proportional to $\gamma_\mu L$ are shown explicitly. The total vertex function $\Gamma_\mu^Z(p,k)$ can therefore be written as

$$\begin{aligned}\Gamma_\mu^Z(p,k) = & \frac{G_F}{\sqrt{2}} \frac{e}{2\pi^2} M_Z^2 \frac{\cos \theta}{\sin \theta} \sum_j V_{js}^* V_{jb} \frac{1}{8} \left\{ \left[-\frac{1}{4-N} + \frac{1}{2} \ln \left(\frac{e^\gamma M_W^2}{4\pi\mu^2} \right) \right] x_j \right. \\ & \left. + \frac{7x_j-8x_j^2+x_j^3}{4(1-x_j)^2} + \frac{4x_j-2x_j^2+x_j^3 \ln x_j}{2(1-x_j)^2} \right\} \gamma_\mu (1-\gamma_5) + \dots, \quad (43)\end{aligned}$$

where the relation $e = g \sin \theta$, $M_W = M_Z \cos \theta$ and $G_F = g^2/(4\sqrt{2}M_W^2)$ have been used.

By comparing the vertex function $\Gamma_\mu^Z(p,k)$ in Eq. (43) with the general expression of the effective coupling $\bar{s}Zb$ [9],

$$\bar{s}Zb = i \sum_j V_{js}^* V_{jb} \frac{G_F}{\sqrt{2}} \frac{e}{2\pi^2} M_Z^2 \frac{\cos \theta}{\sin \theta} C_0(x_j) \bar{s} \gamma_\mu (1-\gamma_5) b, \quad (44)$$

one can extract out the basic function $C_0(x_t)$ in the unitary gauge

$$\begin{aligned}C_0(x_t) = & \frac{1}{8} \left[-\frac{1}{4-N} + \frac{1}{2} \ln \left(\frac{e^\gamma M_W^2}{4\pi\mu^2} \right) \right] x_t \\ & + \frac{7x_t-8x_t^2+x_t^3}{32(1-x_t)^2} + \frac{4x_t-2x_t^2+x_t^3}{16(1-x_t)^2} \ln[x_t], \quad (45)\end{aligned}$$

where $x_t = m_t^2/M_W^2$. Again, a divergent- and a μ -dependent term also appear here!

By combing the function $D_0(x_t)$ in Eq. (25) with the function $C_0(x_t)$ in Eq. (45), we find the function $Z_0(x_t)$ in the unitary gauge

$$\begin{aligned}Z_0(x_t) = & C_0(x_t) + \frac{1}{4} D_0(x_t) \\ = & \frac{108x_t - 259x_t^2 + 163x_t^3 - 18x_t^4}{144(1-x_t)^3} - \frac{8 - 50x_t + 63x_t^2 + 6x_t^3 - 24x_t^4}{72(1-x_t)^4} \ln[x_t], \quad (46)\end{aligned}$$



FIG. 2: Two “tree” Feynman diagrams contributing to $b \rightarrow sl^-l^+$ decays, where the oval denotes the $\bar{s}\gamma b$ vertex and $\bar{s}Zb$ vertex ($k^2 > 0$), respectively.

which is indeed finite, gauge independent and identical with the one as given in Eq. (32) [8, 9]. It is worth of stressing that the divergent- and μ -dependent terms in both $C_0(x_t)$ and $D_0(x_t)$ in the unitary gauge was canceled in the linear combination, and the gauge independence of $Z_0(x_t)$ is true in unitary gauge.

For the semi-leptonic decays $b \rightarrow sl^+l^-$, as illustrated in Fig. 2, it can proceed through the decay chain $b \rightarrow s\gamma^* \rightarrow sl^+l^-$ or $b \rightarrow sZ^* \rightarrow sl^+l^-$, and therefore both effective coupling $\bar{s}\gamma b$ and $\bar{s}Zb$ should contribute simultaneously. The divergent-term, the μ -dependent term and gauge dependence appeared in both $D_0(x_t)$ and $C_0(x_t)$ are canceled each other when one sum up all contributions together, which is consistent with the general expectation.

To summarize, we have recalculated at one-loop level the effective coupling $\bar{s}\gamma b$ and $\bar{s}Zb$ in the unitary gauge and extracted the relevant basic functions $C_0(x_t)$, $D_0(x_t)$ and $D'_0(x_t)$. We start from the calculation of Fig. 1a for the case of $V = \gamma$ and confirmed the analytical results presented in Ref. [5], but our calculations for Fig. 1b are different from the corresponding results as given in Ref. [6, 7]. We finally calculated the effective coupling $\bar{s}Zb$ and made a linear combination of the basic function $D_0(x_t)$ and $C_0(x_t)$ as a means of cross checking. We found that (a) the function $D'_0(x_t)$ is identical with those known results; (b) the divergent- and μ -dependent terms appeared in both $C_0(x_t)$ and $D_0(x_t)$ function in the unitary gauge are canceled each other in a linear combination $Z_0(x_t) = C_0(x_t) + D_0(x_t)/4$; and (c) the resultant $Z_0(x_t)$ function, which governs the semi-leptonic $b \rightarrow sl^+l^-$ decays, is finite, gauge independent and identical with those as given in Ref. [8, 9].

Acknowledgments

We are very grateful to Zhao-hua Xiong, Wen-juan Zou and Lin-xia Lü for helpful discussions. This work is partly supported by the National Natural Science Foundation of China under Grant No.10575052, and by the Specialized Research Fund for the doctoral Program of higher education (SRFDP) under Grant No. 20050319008.

-
- [1] T. Hurth, Rev. Mod. Phys. **75** (2003) 1159.
 - [2] T. Inami and C.S. Lim, Prog.Theor.Phys. **65** (1981) 297; *ibid* **65** (1981) 1772E.
 - [3] N.G. Deshpande and G. Eilam, Phys. Rev. D **26** (1982) 2463.
 - [4] N.G. Deshpande and M. Nazerimonfared, Nucl. Phys. B **213** (1983) 390.
 - [5] Swee-Ping Chia, Phys. Lett. B **130** (1983) 315.
 - [6] Swee-Ping Chia and G. Rajagopal, Phys. Lett. B **156** (1985) 405.
 - [7] Swee-Ping Chia, Phys. Lett. B **240** (1990) 465.
 - [8] G. Buchalla, A.J. Buras, and M.K. Harlander, Nucl. Phys. B **349** (1991)1.
 - [9] G. Buchalla, A.J. Buras, and M.E. Lautenbacher, Rev. Mod. Phys. **68** (1996)1125;
A.J.Buras, in Probing the Standard Model of Particle Interactions, edited by F. David and
R. Gupta Elsevier Science B.V., Amsterdam, 1998; hep-ph/9806471.